CANADIAN JOURNAL OF FOREST RESEARCH

JOURNAL CANADIEN DE RECHERCHE FORESTIÈRE

Volume 19, 1989

Author Index / Index des auteurs

Adams, K.B. see Stergas, R.L., 904

Adams, M.B. see Cunningham, M., 335

Adams, W.T. see Li, P., 149

Adams, W.T. see Kaya, Z., 1124

Adams, W.T. see Erickson, V.J., 1248

Agee, J.K. see Stuart, J.D., 1096

Agnard, J.P. see Gagnon, P.A., 860

Amaranthus, M.P., and Perry, D.A. Interaction effects of vegetation type and Pacific madrone soil inocula on survival, growth, and mycorrhiza formation of Douglas-fir, 550

Amateis, R.L. see Newberry, J.D., 679

Amman, G.D., Thier, R.W., McGregor, M.D., and Schmitz, R.F. Efficacy of verbenone in reducing lodgepole pine infestation by mountain pine beetles in Idaho, 60

Amman, G.D., McGregor, M.D., Schmitz, R.F., and Oakes, R.D. Erratum: Susceptibility of lodgepole pine to infestation by mountain pine beetles following partial cutting of stands, 284

Amman, G.D. see Schmitz, R.F., 566

Amrhein, J.F. see Fairweather, S.E., 1169

Anderson, D.H. A mathematical model for fire containment, 997

Anderson, R.L. see Foster, G.S., 534

Arbaugh, M.J. see Peterson, D.L., 401

Arbaugh, M.J. see Peterson, D.L., 530

Arnebrant, K. see Nohrstedt, H.-Ö., 323

Arnott, J.T. see O'Reilly, C., 122

Arthur, L.M. see Van Kooten, G.C., 463

Arvanitis, L.G. see Reich, R.M., 1550

Aussenac, G. see Kaushal, P., 1351

Baath, E. see Nohrstedt, H.-Ö., 323

Baes, C.F., III. see Bondietti, E.A., 586

Baker, W.L. Effect of scale and spatial heterogeneity on fireinterval distributions, 700

Baldwin, V.C., Jr., Feduccia, D.P., and Haywood, J.D. Postthinning growth and yield of row-thinned and selectively thinned loblolly and slash pine plantations, 247

Banerjee, S. see Roy, D.N., 842

Bare, B.B. see O'Hara, A.J., 715

Barker, J. see Weetman, G.F., 1501

Barker, J. see Weetman, G.F., 1512

Bartolome, J.W. see McClaran, M.P., 580

Bassman, J.H. Influence of two site preparation treatments on ecophysiology of planted *Picea engelmannii* × *glauca* seedlings, 1359

Baxter, R., Brown, S.N., England, N.F., Ludlow, C.H.M., Taylor, S.L., and Womack, R.W. Production of clonal plantlets of tropical pine in tissue culture via axillary shoot activation, 1338

Beadle, C.L. see West, P.W., 270

Beadle, C.L. see Hallam, P.M., 1235

Beck, D.E. see Bowling, E.H., 405

Beck, J.A., and Trevitt, A.C.F. Forecasting diurnal variations in meteorological parameters for predicting fire behaviour, 791

Becker, M. The role of climate on present and past vitality of silver fir forests in the Vosges mountains of northeastern France, 1110

Bekkaoui, F., and Dunstan, D.I. Permeabilization of *Picea glauca* protoplasts to macromolecules, 1316

Bell, R.L., and Binkley, D. Soil nitrogen mineralization and immobilization in response to periodic prescribed fire in a loblolly pine plantation, 816

Bella, I.E. see Cieszewski, C.J., 1151

Bennett, J.P. see Berrang, P., 519

Bentley, C.V. see Fayle, D.C.F., 440

Bentzer, B.G., Foster, G.S., Hellberg, A.R., and Podzorski, A.C. Trends in genetic and environmental parameters, genetic correlations, and response to indirect selection for 10-year volume in a Norway spruce clonal experiment, 897

Bentzer, B.G. see Foster, G.S., 806

Bergdahl, D.R. see Mazzola, M., 88

Bergeron, Y. see Bouchard, A., 1146

Bergeron, Y. see Harvey, B.D., 1458

Bernier, B. see Paré, D., 24

Bernier, B. see Paré, D., 132

Bernier, B. see Paré, D., 135

Berrang, P., Karnosky, D.F., and Bennett, J.P. Natural selection for ozone tolerance in *Populus tremuloides*: field verification, 519

Bethune, T. see Dunstan, D.I., 1299

Bevilacqua, E. see Martell, D.L., 1555

Bidlake, W.R., and Black, R.A. Vertical distribution of leaf area in *Larix occidentalis*: a comparison of two estimation methods, 1131

Binder, W. see Vidaver, W., 1478

Binkley, D. see Strader, R.H., 798

Binkley, D. see Bell, R.L., 816

Binkley, D. see France, E.A., 1648

Black, R.A. see Bidlake, W.R., 1131 Blanchette, R.A. see Enebak, S.A., 225

Blanchette, R.A. see Burnes, T.A., 1642

Blenis, P.V., Wiggins, K.L., Cunningham, J.E., and Pickard, M.A. Erratum: Maltol protects against infection of lodgepole pine seedlings by western gall rust, 683

Blenis, P.V., Mugala, M.S., and Hiratsuka, Y. Soil affects Armillaria root rot of lodgepole pine, 1638

Blenis, P.V. see Mugala, M.S., 685

Bockheim, J.G., Leide, J.E., and Frelich, L.E. Red pine growth and chemical composition of foliage and forest floors across a precipitation-chemistry gradient in Wisconsin, 1543

Bockheim, J.G. see Frelich, L.E., 113

Bondietti, E.A., Baes, C.F., III, and McLaughlin, S.B. Radial trends in cation ratios in tree rings as indicators of the impact of atmospheric deposition on forests, 586

Bongarten, B.C. see Surles, S.E., 471

Borden, J.H. see Lindgren, B.S., 65

Borden, J.H. see Hunt, D.W.A., 1275

Borders, B.E., and Shiver, B.D. Herbicide field studies in forestry: statistical and other considerations, 768

Borghetti, M., Cocco, S., Lambardi, M., and Raddi, S. Response to water stress of Italian alder seedlings from diverse geographic origins, 1071

Bouchard, A., Dyrda, S., Bergeron, Y., and Meilleur, A. The use of notary deeds to estimate the changes in the composition of 19th century forests, in Haut-Saint-Laurent, Quebec, 1146

Bowden, R.D., Geballe, G.T., and Bowden, W.B. Foliar uptake of ¹⁵N from simulated cloud water by red spruce (*Picea rubens*) seedlings, 382

Bowden, W.B. see Bowden, R.D., 382

Bower, R.C., Ross, S.D., and Dunsworth, B.G. Effect of GA₃ treatment timing in relation to natural day length on flowering and sex expression in *Chamaecyparis nootkatensis*, 1422

Bowling, E.H., Burkhart, H.E., Burk, T.E., and Beck, D.E. A stand-level multispecies growth model for Appalachian hard-woods, 405

Brix, H. see Jozsa, L.A., 1137

Brooke, R.C. see Vidaver, W., 1478

Brown, S.N. see Baxter, R., 1338

Buford, M.A. Mean stem size and total volume development of various loblolly pine seed sources planted at one location, 396

Bullard, S.H. see Cubbage, F.W., 759
Burdon, R.D. Farly selection in tree brees

Burdon, R.D. Early selection in tree breeding: principles for applying index selection and inferring input parameters, 499

Burdon, R.D. Erratum: Early selection in tree breeding: principles for applying index selection and inferring input parameters, 1210

Burgess, D. see Hendrickson, O.Q., 725

Burgess, D. see Hendrickson, O.Q., 936

Burk, T.E. see Bowling, E.H., 405

Burkhart, H.E. see Bowling, E.H., 405

Burkhart, H.E. see Newberry, J.D., 679

Burnes, T.A., Blanchette, R.A., Stewart, W.K., and Mohn, C.A. Screening jack pine seedlings for resistance to Cronartium quercuum f.sp. banksianae and Endocronartium harknessii, 1642

Burns, S.F. see Schaetzl, R.J., 1

Bywaters, T. see Dell, B., 427

Cai, M. see Stoehr, M.U., 1634

Campagna, M.A., and Margolis, H.A. Influence of short-term atmospheric CO₂ enrichment on growth, allocation patterns, and biochemistry of black spruce seedlings at different stages of development, 773

Campbell, R.K., Pawuk, W.A., and Harris, A.S. Microgeographic genetic variation of Sitka spruce in southeastern Alaska, 1004

Campbell, R.K. see Kaya, Z., 1124

Cannon, G.P. see Struve, D.K., 505

Cannon, W.N., Jr. see Roberts, B.R., 1200

Carlyle, J.C., Turvey, N.D., Hopmans, P., and Downes, G.M. Stem deformation in *Pinus radiata* associated with previous land use, 96

Carmean, W.H., and Lenthall, D.J. Height-growth and site-index curves for jack pine in north central Ontario, 215

Caron, G.E., and Powell, G.R. Cone size and seed yield in young Picea mariana trees, 351

Caron, G.E., and Powell, G.R. Patterns of seed-cone and pollencone production in young *Picea mariana* trees, 359

Castello, J.D. see Nelson, C.L., 389

Caulfield, J.P., Shoulders, E., and Lockaby, B.G. Risk-efficient species-site selection decisions for southern pines, 743

Charles, D.A. see Roy, D.N., 842

Chatarpaul, L. see Hendrickson, O.Q., 725

Chong, L.J. see Lindgren, B.S., 65

Chrosciewicz, Z. Prediction of forest-floor moisture content on jack pine cutovers, 239

Chrosciewicz, Z. Prediction of forest-floor moisture content under diverse jack pine canopy conditions, 1483

Cieszewski, C.J., and Bella, I.E. Polymorphic height and site index curves for lodgepole pine in Alberta, 1151 Cocco, S. see Borghetti, M., 1071

Cohen, D.H., and Sinclair, S.A. An inventory of innovative technology use in North American processing of wood structural panels and softwood lumber, 1629

Colangeli, A.M., and Owens, J.N. Postdormancy seed-cone development and the pollination mechanism in western hemlock (*Tsuga heterophylla*), 44

Colangeli, A.M. see Owens, J.N., 1051

Colombo, S.J. see Odlum, K.D., 274

Comeau, P.G., and Kimmins, J.P. Above- and below-ground biomass and production of lodgepole pine on sites with differing soil moisture regimes, 447

Connor, K.F., and Lanner, R.M. Age-related changes in shoot growth components of Great Basin bristlecone pine, 933

Cook, E.R. see Innes, J.L., 1174

Corbin, J.P. see Prescott, C.E., 309

Corbin, J.P. see Prescott, C.E., 489

Côté, B., and Dawson, J.O. Effects of temperature regime and fertilization on nitrogenase activity of black alder seedlings during autumn in Illinois, U.S.A., 1644

Criddle, R.S. see Hansen, L.D., 606

Cronan, C.S. see Rustad, L.E., 18

Cubbage, F.W., Wojtkowski, P.A., and Bullard, S.H. Crosssectional estimation of empirical southern United States pulpwood harvesting cost functions, 759

Cunia, T. see Gillespie, A.J.R., 664

Cunningham, J.E. see Blenis, P.V., 683

Cunningham, M., Adams, M.B., Luxmoore, R.J., Post, W.M., and DeAngelis, D.L. Quick estimates of root length, using a video image analyzer, 335

Cushon, G.H., and Feller, M.C. Asymbiotic nitrogen fixation and denitrification in a mature forest in coastal British Columbia, 1194

Cushon, G.H. see Lindgren, B.S., 65

Czaplewski, R.L. Graphical analysis of stem taper in model building, 522

Dale, V.H., and Franklin, J.F. Potential effects of climate change on stand development in the Pacific Northwest, 1581

Dang, Q.L., and Lieffers, V.J. Assessment of patterns of response of tree ring growth of black spruce following peatland drainage, 924

Danielson, R.M., and Visser, S. Host response to inoculation and behaviour of introduced and indigenous ectomycorrhizal fungi of jack pine grown on oil-sands tailings, 1412

Davis, D.D. see Long, R.P., 1603

Davis, J.M., and Keathley, D.E. Detection and analysis of T-DNA in crown gall tumors and kanamycin-resistant callus of Robinia pseudoacacia, 1118

Dawson, J.O. see Neave, I.A., 12

Dawson, J.O. see Côté, B., 1644

Dealy, J.E. see Farr, W.A., 1203

DeAngelis, D.L. see Cunningham, M., 335

DeHayes, D.H., Ingle, M.A., and Waite, C.E. Nitrogen fertilization enhances cold tolerance of red spruce seedlings, 1037

Dell, B., and Bywaters, T. Copper deficiency in young Eucalyptus maculata plantations, 427

DeLucia, E.H. see Neave, I.A., 12

DeMars, D.J. see Farr, W.A., 1203

Désaulniers, G. see Gervais, C., 595

De Vries, P.G., and Schreuder, H.T. Estimation of stand means and other ratios in point sampling, 1067

Dixon, R.K. see Yin, X., 204

Dobesberger, E.J. A sequential decision plan for the management of the eastern hemlock looper, *Lambdina fiscellaria fiscellaria* (Lepidoptera: Geometridae), in Newfoundland, 911

Downes, G.M. see Carlyle, J.C., 96

Dumroese, R.K. see Stiff, C.M., 1330

Dunstan, D.I. Introduction, 1297

Dunstan, D.I., Bethune, T., Kikcio, S., and Lapp, M.S. In vitro shoot formation among 17-year-old Douglas-fir provenances, 1299

Dunstan, D.I. see Bekkaoui, F., 1316

Dunsworth, B.G. see Bower, R.C., 1422

Duryea, M.L. see Schuch, U.K., 192

Duryea, M.L. see Schuch, U.K., 198

Dyrda, S. see Bouchard, A., 1146 Dyrness, C.T., Van Cleve, K., and Levison, J.

Dyrness, C.T., Van Cleve, K., and Levison, J.D. The effect of wildfire on soil chemistry in four forest types in interior Alaska, 1389

Eatough, D.J. see Hansen, L.D., 606

Edmonds, R.L., and Eglitis, A. The role of the Douglas-fir beetle and wood borers in the decomposition of and nutrient release from Douglas-fir logs, 853

Eglitis, A. see Edmonds, R.L., 853

Ellis, D.D., Lazaroff, W.R., Roberts, D.R., Flinn, B.S., and Webb, D.T. The effect of antibiotics on elongation and callus and bud formation from embryonic tissue of *Picea glauca*, 1343

Ellis, D.D. see Webb, D.T., 1303
Enebak, S.A., and Blanchette, R.A. Canker formation and decay in sugar maple and paper birch infected by Cerrena unicolor, 225

England, N.F. see Baxter, R., 1338

Erickson, V.J., and Adams, W.T. Mating success in a coastal Douglas-fir seed orchard as affected by distance and floral phenology, 1248

Faaland, B.H. see O'Hara, A.J., 715

Fairweather, S.E., and Amrhein, J.F. A test of a nonparametric distance-based estimator for seedling density, 1169

Farr, W.A., DeMars, D.J., and Dealy, J.E. Height and crown width related to diameter for open-grown western hemlock and Sitka spruce, 1203

Fayle, D.C.F., and Bentley, C.V. Temporal changes in growthlayer patterns of plantation-grown red oak and red pine, 440

Feduccia, D.P. see Baldwin, V.C., Jr., 247

Feller, M.C. see Cushon, G.H., 1194

Feng, C.C. see Feng, J.C., 378

Feng, J.C., Feng, C.C., and Sidhu, S.S. Determination of hexazinone residue and its release from a granular formulation under forest conditions, 378

Feret, P.P. see Johnsen, K.H., 1598

Fielder, P., and Owens, J.N. A comparative study of shoot and root development of interior and coastal Douglas-fir seedlings, 539

Fielder, P., and Owens, J.N. Erratum: A comparative study of shoot and root development of interior and coastal Douglas-fir seedlings, 1210

Finney, M.A., and Martin, R.E. Fire history in a Sequoia sempervirens forest at Salt Point State Park, California, 1451

Firestone, M.K. see Hart, S.C., 185

Fisher, R.F. see Verbyla, D.L., 179

Flannigan, M.D., and Wotton, B.M. A study of interpolation methods for forest fire danger rating in Canada, 1059

Flinn, B.S. see Webb, D.T., 1303

Flinn, B.S. see Ellis, D.D., 1343

Fortin, J.A. see Jobidon, R., 1291

Foster, G.S., and Anderson, R.L. Indirect selection and clonal propagation of loblolly pine seedlings enhance resistance to fusiform rust, 534

Foster, G.S., Bentzer, B.G., Hellberg, A.R., and Podzorski, A.C. Heights and growth habit of Norway spruce rooted cuttings compared between two serial propagation cycles, 806

Foster, G.S. see Bentzer, B.G., 897

Fournier, R. see Weetman, G.F., 1501

Fournier, R. see Weetman, G.F., 1512

Fowler, D.P. see Hansen, L.D., 606

Frampton, L.J., Jr., and Jett, J.B. Juvenile wood specific gravity of loblolly pine tissue culture plantlets and seedlings, 1347

France, E.A., Binkley, D., and Valentine, D. Soil chemistry changes after 27 years under four tree species in southern Ontario, 1648

Franklin, J.F. see Dale, V.H., 1581

Freedman, B. see Morash, R., 347

Frelich, L.E., Bockheim, J.G., and Leide, J.E. Historical trends in tree-ring growth and chemistry across an air-quality gradient in Wisconsin, 113

Frelich, L.E. see Lorimer, C.G., 651

Frelich, L.E. see Bockheim, J.G., 1543

Froelich, R.C. Annual variation in fusiform rust infection of slash and loblolly pine seed lots in time-replicated plantings, 1531

Fuchigami, L.H. see Schuch, U.K., 192 Fuchigami, L.H. see Schuch, U.K., 198

Futai, K., and Sutherland, J.R. Pathogenicity and attraction to host extracts of Canadian pinewood nematodes: studies with Scots pine, western larch, and black spruce seedlings, 1256

Gagnon, P.A., and Agnard, J.P. Twin-camera fixed-base photography of tree plots: possibilities and accuracy of a system, 860

Gara, R.I. see Stuart, J.D., 1096

Gassmann, H.I. Optimal harvest of a forest in the presence of uncertainty, 1267

Geballe, G.T. see Bowden, R.D., 382

Germain, A. see Weetman, G.F., 1501

Gervais, C., Roy, G., Grandtner, M.M., and Désaulniers, G. The B chromosomes of *Claytonia caroliniana* (Portulacaceae) and maple forest dieback, 595

Gilbert, J. see Punter, D., 865

Gillespie, A.J.R., and Cunia, T. Linear regression models for biomass table construction, using cluster samples, 664

Gilliam, F.S., Sigmon, J.T., Reiter, M.A., and Krovetz, D.O. Elevational and spatial variation in daytime ozone concentrations in the Virginia Blue Ridge Mountains: implications for forest exposure, 422

Gilliam, F.S. see Sigmon, J.T., 1240

Gonzalez, J.S. Circumferential variation in the relative density of lodgepole pine, 276

Gower, S.T., and Grier, C.C. Aboveground organic matter and production of a montane forest on the eastern slopes of the Washington Cascade Range, 515

Grandtner, M.M. see Gervais, C., 595

Greulich, F.E. The calculation of average yarding distance to a centralized landing, given a polygonal mesh approximation of the setting surface, 141

Grier, C.C. see Gower, S.T., 515

Gries, G. see Hunt, D.W.A., 1275

Grossnickle, S.C. Shoot phenology and water relations of *Picea glauca*, 1287

Guehl, J.M. see Kaushal, P., 1351

Guries, R.P. see Rudolph, T.D., 707

Hadley, K.S. see Veblen, T.T., 1218

Haines, B.L. see Montagnini, F., 1226

Hall, J.P. A proposal for the development of a land race of Sitka spruce in Newfoundland, 939

Hallam, P.M., and Reid, J.B. Seasonal and genetic variation in frost hardiness of Eucalyptus delegatensis, 480

Hallam, P.M., Reid, J.B., and Beadle, C.L. Frost hardiness of commercial Eucalyptus species at different elevations, 1235

Hamrick, J.L. see Surles, S.E., 471

Hann, D.W. see Maguire, D.A., 557

Hansen, L.D., Lewis, E.A., Eatough, D.J., Fowler, D.P., and Criddle, R.S. Prediction of long-term growth rates of larch clones by calorimetric measurement of metabolic heat rates, 606

Hanson, P.J. see Norby, R.J., 889

Harmon, M.E. see Kelsey, R.G., 1030

Harr, R.D. see Martin, C.W., 35

Harris, A.S. see Campbell, R.K., 1004

Hart, S.C., and Firestone, M.K. Evaluation of three in situ soil nitrogen availability assays, 185

Harvey, B.D., and Bergeron, Y. Site patterns of natural regeneration following clear-cutting in northwestern Quebec, 1458

Haywood, J.D. see Baldwin, V.C., Jr., 247

Heagle, A.S. see Shafer, S.R., 821

Heaman, J.C. see Woods, J.H., 54

Heilmann, P.E. see Rogers, D.L., 372

Hellberg, A.R. see Foster, G.S., 806

Hellberg, A.R. see Bentzer, B.G., 897

Hendrickson, O.Q., Chatarpaul, L., and Burgess, D. Nutrient cycling following whole-tree and conventional harvest in northern mixed forest, 725

Hendrickson, O.Q., and Burgess, D. Nitrogen-fixing plants in a cut-over lodgepole pine stand of southern British Columbia, 936

Herring, L. see Ying, C.C., 832

Higgins, C.J. see Lindgren, B.S., 65

Hiratsuka, Y. see Mugala, M.S., 685

Hiratsuka, Y. see Blenis, P.V., 1638

Holland, L. see Horgan, K., 1309

Holopainen, T., and Nygren, P. Effects of potassium deficiency and simulated acid rain, alone and in combination, on the ultrastructure of Scots pine needles, 1402

Holthausen, D.M. see Hyberg, B.T., 1014

Honer, T.W. see Messier, C., 1470

Hopmans, P. see Carlyle, J.C., 96

Horgan, K., and Holland, L. Rooting micropropagated shoots from mature radiata pine, 1309

Host, C.E. see Zak, D.R., 1521

Howard, A.F. A sequential approach to sampling design for time studies of cable yarding operations, 973

Howard, A.F., and Thérien, G. Regression analysis using auxiliary information from time studies of cable yarding operations, 1262

Hunt, D.W.A., Borden, J.H., Lindgren, B.S., and Gries, G. The role of autoxidation of α-pinene in the production of pheromones of *Dendroctonus ponderosae* (Coleoptera: Scolytidae), 1275

Hyberg, B.T., and Holthausen, D.M. The behavior of nonindustrial private forest landowners, 1014

Ingle, M.A. see DeHayes, D.H., 1037

Innes, J.L., and Cook, E.R. Tree-ring analysis as an aid to evaluating the effects of pollution on tree growth, 1174

Jett, J.B. see Struve, D.K., 505

Jett, J.B. see Frampton, L.J., Jr., 1347

Jobidon, R., Thibault, J.R., and Fortin, J.A. Effect of straw

residues on black spruce seedling growth and mineral nutrition, under greenhouse conditions, 1291

Johnsen, K.H., Feret, P.P., and Seiler, J.R. Root growth potential and bud dormancy of 2+0 eastern white pine grown in a Virginia nursery, 1598

Johnson, A.F., Woodard, P.M., and Titus, S.J. Lodgepole pine and white spruce crown fuel weights predicted from height and crown width, 527

Johnson, D.L. see Schaetzl, R.J., 1

Johnson, W.C. see Schiffman, P.M., 69

Jones, E.A. see Reed, D.D., 948

Jozsa, L.A., and Brix, H. The effects of fertilization and thinning on wood quality of a 24-year-old Douglas-fir stand, 1137

Kalisz, P.J. see Stringer, J.W., 627

Kalra, Y.P., Maynard, D.G., and Radford, F.G. Microwave digestion of tree foliage for multi-element analysis, 981

Kang, H.C. see Rudolph, T.D., 707

Karnosky, D.F. see Berrang, P., 519

Karnosky, D.F. see McLaughlin, J., 1334

Kauffman, J.B., and Martin, R.E. Fire behavior, fuel consumption, and forest-floor changes following prescribed understory fires in Sierra Nevada mixed conifer forests, 455

Kaushal, P., Guehl, J.M., and Aussenac, G. Differential growth response to atmospheric carbon dioxide enrichment in seedlings of Cedrus atlantica and Pinus nigra ssp. Laricio var. Corsicana, 1351

Kaya, Z., Campbell, R.K., and Adams, W.T. Correlated responses of height increment and components of increment in 2-year-old Douglas-fir, 1124

Keathley, D.E. see Davis, J.M., 1118

Kelsey, R.G., and Harmon, M.E. Distribution and variation of extractable total phenols and tannins in the logs of four conifers after 1 year on the ground, 1030

Kikcio, S. see Dunstan, D.I., 1299

Kimmins, J.P. see Comeau, P.G., 447

Kimmins, J.P. see Messier, C., 1470

Kiss, G.K. Engelmann \times Sitka spruce hydrids in central British Columbia, 1190

Klein, J.I. Effects of incomplete block design and stand structure on heritability of growth in a jack pine family test in Manitoba, 1044

Klein, R.M., Perkins, T.D., and Myers, H.L. Nutrient status and winter hardiness of red spruce foliage, 754

Klein, R.M. see Liedeker, H., 145

Kleiner, K.W., Montgomery, M.E., and Schultz, J.C. Variation in leaf quality of two oak species: implications for stand susceptibility to gypsy moth defoliation, 1445

Knowles, P. see Perry, D.J., 509

Konar, S.K. see Roy, D.N., 842

Koran, Z. see Nlombi, K., 986

Kourtz, P. Two dynamic programming algorithms for forest fire resource dispatching, 106

Kraft, J.M. see Radwan, M.A., 1429

Kremer, A., and Xu, L. Relationship between first-season free growth components and later field height growth in maritime pine (*Pinus pinaster*), 690

Kriebel, H.B. Genetic improvement of sugar maple for high sap sugar content. I. Clone selection and seed orchard development, 917

Krovetz, D.O. see Gilliam, F.S., 422

Kwon, K.W., and Pallardy, S.G. Temporal changes in tissue water relations of seedlings of *Quercus acutissima*, Q. alba, and Q. stellata subjected to chronic water stress, 622

Lambardi, M. see Borghetti, M., 1071

Lanner, R.M. see Connor, K.F., 933

Lapp, M.S. see von Rudloff, E., 848

Lapp, M.S. see Dunstan, D.I., 1299

Lardner, M.A. see Peterson, D.L., 401 Lazaroff, W.R. see Ellis, D.D., 1343

Leide, J.E. see Frelich, L.E., 113

Leide, J.E. see Bockheim, J.G., 1543

Lenthall, D.J. see Carmean, W.H., 215

Levison, J.D. see Dyrness, C.T., 1389

Lewis, E.A. see Hansen, L.D., 606

Li, P., and Adams, W.T. Range-wide patterns of allozyme variation in Douglas-fir (Pseudotsuga menziesii), 149

Liedeker, H., and Klein, R.M. Characteristic discolorations of coniferous foliage related to spruce decline in Europe and North America, 145

Lieffers, V.J. see Dang, Q.L., 924

Lindgren, B.S., Borden, J.H., Cushon, G.H., Chong, L.J., and Higgins, C.J. Reduction of mountain pine beetle (Coleoptera: Scolytidae) attacks by verbenone in lodgepole pine stands in British Columbia, 65

Lindgren, B.S. see Hunt, D.W.A., 1275

Lister, G.R. see Vidaver, W., 1478

Liu, C.J. A form-class tree-grading algorithm, 137

Lockaby, B.G. see Caulfield, J.P., 743

Long, J.N., and Smith, F.W. Estimating leaf area of Abies lasiocarpa across ranges of stand density and site quality, 930

Long, R.P., and Davis, D.D. Major and trace element concentrations in surface organic layers, mineral soil, and white oak xylem downwind from a coal-fired power plant, 1603

Lorimer, C.G., and Frelich, L.E. A methodology for estimating canopy disturbance frequency and intensity in dense temperate forests, 651

Ludlow, C.H.M. see Baxter, R., 1338

Luxmoore, R.J. see Cunningham, M., 335

MacLean, D.A., and Ostaff, D.P. Patterns of balsam fir mortality caused by an uncontrolled spruce budworm outbreak, 1087

MacLean, D.A. see Ostaff, D.P., 1077

Magnussen, S., and Yeatman, C.W. Height growth components in inter- and intra-provenance jack pine families, 962

Maguire, D.A., and Hann, D.W. The relationship between gross crown dimensions and sapwood area at crown base in Douglasfir, 557

Mallett, K.I. see Mugala, M.S., 685

Mallik, A.U. see Thompson, I.D., 524

Manion, P.D. see Nelson, C.L., 389

Margolis, H.A. see Pothier, D., 432

Margolis, H.A. see Campagna, M.A., 773

Margolis, H.A. see Pothier, D., 1564

Martel, D.L., Bevilacqua, E., and Stocks, B.J. Modelling seasonal variation in daily people-caused forest fire occurrence, 1555

Martin, C.W., and Harr, R.D. Logging of mature Douglas-fir in western Oregon has little effect on nutrient output budgets, 35

Martin, R.E. see Kauffman, J.B., 455

Martin, R.E. see Finney, M.A., 1451

Maynard, D.G. see Kalra, Y.P., 981

Mazzola, M., and Bergdahl, D.R. The phenology and spore dissemination patterns of *Uredinopsis mirabilis* on *Abies* balsamea and *Onoclea sensibilis*, 88

McAlpine, R.S. Temporal variations in elliptical forest fire shapes, 1496

McClaran, M.P., and Bartolome, J.W. Fire-related recruitment

in stagnant Quercus douglasii populations, 580

McClenahen, J.R., Vimmerstedt, J.P., and Scherzer, A.J. Elemental concentrations in tree rings by PIXE: statistical variability, mobility, and effects of altered soil chemistry, 880

McDermott, J.M., and Robinson, R.A. Provenance variation for disease resistance in *Pseudotsuga menziesii* to the Swiss needlecast pathogen, *Phaeocryptopus gaeumannii*, 244

McGregor, M.D. see Amman, G.D., 60

McGregor, M.D. see Amman, G.D., 284

McGregor, M.D. see Schmitz, R.F., 566

McKeand, S.E. see Struve, D.K., 505

McKittrick, R.C. see Stanturf, J.A., 279

McLaughlin, J., and Karnosky, D.F. Controlling vitrification in Larix decidua via culture media manipulation, 1334

McLaughlin, S.B. see Bondietti, E.A., 586

McMinn, R.G. see Thomson, A.J., 257

McMinn, R.G. see Thomson, A.J., 262

Means, J.E. Estimating the date of a single bole scar by counting tree rings in increment cores, 1491

Meilleur, A. see Bouchard, A., 1146

Merkle, S.A., and Wiecko, A.T. Regeneration of *Robinia* pseudoacacia via somatic embryogenesis, 285

Messier, C., Honer, T.W., and Kimmins, J.P. Photosynthetic photon flux density, red:far-red ratio, and minimum light requirement for survival of Gaultheria shallon in western red cedar - western hemlock stands in coastal British Columbia, 1470

Millar, C.I. Allozyme variation of bishop pine associated with pygmy-forest soils in northern California, 870

Mohammed, G.H., Patel, K.R., and Vidaver, W.E. The control of adventitious root production in tissue-cultured Douglas-fir, 1322

Mohn, C.A. see Burnes, T.A., 1642

Montagnini, F., Haines, B.L., Swank, W.T., and Waide, J.B. Nitrification in undisturbed mixed hardwoods and manipulated forests in the southern Appalachian Mountains of North Carolina, U.S.A., 1226

Montgomery, M.E. see Kleiner, K.W., 1445

Moore, J.A. see Real, P.L., 1538

Morash, R., and Freedman, B. The effects of several herbicides on the germination of seeds in the forest floor, 347

Mosseler, A. Interspecific pollen-pistil incongruity in Salix, 1161
Mugala, M.S., Blenis, P.V., Hiratsuka, Y., and Mallett, K.I.
Infection of lodgepole pine and white spruce by Alberta isolates of Armillaria, 685

Mugala, M.S. see Blenis, P.V., 1638

Munson, A.D., and Timmer, V.R. Site-specific growth and nutrition of planted *Picea mariana* in the Ontario Clay Belt. I. Early performance, 162

Munson, A.D., and Timmer, V.R. Site-specific growth and nutrition of planted *Picea mariana* in the Ontario Clay Belt. II. Effects of nitrogen fertilization, 171

Murray, M.D. see Radwan, M.A., 1429

Myers, H.L. see Klein, R.M., 754

Myers, R.K. see Zahner, R., 612

Neave, I.A., Dawson, J.O., and DeLucia, E.H. Autumnal photosynthesis is extended in nitrogen-fixing European black alder compared with white basswood: possible adaptive significance, 12

Nelson, C.L., Castello, J.D., and Manion, P.D. Preliminary investigation of the use of indirect immunofluorescence to detect conidia of *Brunchorstia pinea*, 389

Newberry, J.D., Burkhart, H.E., and Amateis, R.L. Individual tree merchantable volume to total volume ratios based on

geometric solids, 679

Newberry, J.D. see Real, P.L., 1538

Newton, M. see White, D.E., 232

Nicholas, N.S., and Zedaker, S.M. Ice damage in spruce-fir forests of the Black Mountains, North Carolina, 1487

Nlombi, K., et Koran, Z. Les caractéristiques papetières du bois attaqué par la tordeuse des bourgeons de l'épinette, 986

Nohrstedt, H.-Ö., Arnebrant, K., Bååth, E., and Söderström, B. Changes in carbon content, respiration rate, ATP content, and microbial biomass in nitrogen-fertilized pine forest soils in Sweden, 323

Norby, R.J., Weerasuriya, Y., and Hanson, P.J. Induction of nitrate reductase activity in red spruce needles by NO₂ and

HNO₃ vapor, 889

Norton, D.A. Tree windthrow and forest soil turnover, 386

Nygren, P. see Holopainen, T., 1402

Oakes, R.D. see Amman, G.D., 284

Oakes, R.D. see Schmitz, R.F., 566

Odlum, K.D., and Colombo, S.J. The influence of night temperature under declining photoperiod on bud initiation in black spruce seedlings, 274

O'Hara, A.J., Faaland, B.H., and Bare, B.B. Spatially constrained timber harvest scheduling, 715

Oppenheimer, M.J., Shiver, B.D., and Rheney, J.W. Ten-year growth response of midrotation slash pine plantations to control of competing vegetation, 329

O'Reilly, C., and Owens, J.N. Polycyclic growth and branching in the upper crown in provenances of lodgepole pine, 79

O'Reilly, C., Arnott, J.T., and Owens, J.N. Effects of photoperiod and moisture availability on shoot growth, seedling morphology, and cuticle and epicuticular wax features of container-grown western hemlock seedlings, 122

O'Reilly, C., and Owens, J.N. Shoot, needle, and cambial growth phenology and branch tracheid dimensions in provenances of

lodgepole pine, 599

Osawa, A. Causality in mortality patterns of spruce trees during a spruce budworm outbreak, 632

Ostaff, D.P., and MacLean, D.A. Spruce budworm populations, defoliation, and changes in stand condition during an uncontrolled spruce budworm outbreak on Cape Breton Island, Nova Scotia, 1077

Ostaff, D.P. see MacLean, D.A., 1087

Ostrofsky, A. see Schwintzer, C.R., 1105

Ouellet, D., et Zarnovican, R. La cond

Ouellet, D., et Zarnovican, R. La conduite des jeunes peuplements de bouleaux jaunes (*Betulla alleghaniensis* Britton): caractéristiques morphologiques, 992

Owens, J.N., and Colangeli, A.M. Promotion of flowering in western hemlock by gibberellin A_{4/7} applied at different stages of bud and shoot development, 1051

Owens, J.N. see Colangeli, A.M., 44

Owens, J.N. see O'Reilly, C., 79

Owens, J.N. see O'Reilly, C., 122

Owens, J.N. see Fielder, P., 539

Owens, J.N. see O'Reilly, C., 599

Owens, J.N. see Fielder, P., 1210

Pallardy, S.G. see Kwon, K.W., 622

Paré, D., and Bernier, B. Origin of the phosphorus deficiency observed in declining sugar maple stands in the Quebec Appalachians, 24

Paré, D., and Bernier, B. Phosphorus-fixing potential of Ah and H horizons subjected to acidification, 132

Paré, D., and Bernier, B. Changes in phosphorus nutrition of sugar maple along a topographic gradient in the Quebec Appalachians, 135

Parkinson, D. see Prescott, C.E., 309

Parkinson, D. see Prescott, C.E., 489

Parkinson, D. see Taylor, B.R., 674

Parkinson, D. see Visser, S., 955

Parsons, W.F.J. see Taylor, B.R., 674

Partin, M.E. see Sigmon, J.T., 1240

Patel, K.R. see Mohammed, G.H., 1322

Pawuk, W.A. see Campbell, R.K., 1004

Payette, S. see Sirois, L., 1571

Penner, M. Optimal design with variable cost and precision requirements, 1591

Perkins, T.D. see Klein, R.M., 754

Perry, D.A. see Amaranthus, M.P., 550

Perry, D.J., and Knowles, P. Allozyme variation in sugar maple at the northern limit of its range in Ontario, Canada, 509

Perry, J.A. see Yin, X., 204

Peterson, D.L., Arbaugh, M.J., and Lardner, M.A. Leaf area of lodgepole pine and whitebark pine in a subalpine Sierra Nevada forest, 401

Peterson, D.L., and Arbaugh, M.J. Estimating postfire survival of Douglas-fir in the Cascade Range, 530

Pickard, M.A. see Blenis, P.V., 683

Piene, H. Spruce budworm defoliation and growth loss in young balsam fir: defoliation in spaced and unspaced stands and individual tree survival, 1211

Piene, H. Spruce budworm defoliation and growth loss in young balsam fir: recovery of growth in spaced stands, 1616

Plamondon, A.P. see Prévost, M., 575

Podzorski, A.C. see Foster, G.S., 806

Podzorski, A.C. see Bentzer, B.G., 897

Poliquin, J. see Pothier, D., 1564

Post, W.M. see Cunningham, M., 335

Pothier, D., Margolis, H.A., and Waring, R.H. Patterns of change of saturated sapwood permeability and sapwood conductance with stand development, 432

Pothier, D., Margolis, H.A., Poliquin, J., and Waring, R.H. Relation between the permeability and the anatomy of jack pine sapwood with stand development, 1564

Powell, G.R. see Caron, G.E., 351

Powell, G.R. see Caron, G.E., 359

Prasad, R. see Roy, D.N., 842

Pregitzer, K.S. see Zak, D.R., 1521

Prescott, C.E., Corbin, J.P., and Parkinson, D. Biomass, productivity, and nutrient-use efficiency of aboveground vegetation in four Rocky Mountain coniferous forests, 309

Prescott, C.E., Corbin, J.P., and Parkinson, D. Input, accumulation, and residence times of carbon, nitrogen, and phosphorus in four Rocky Mountain coniferous forests, 489

Prévost, M., Stein, J., and Plamondon, A.P. Water balance and irrigation planning in a forest tree nursery, 575

Punter, D., and Gilbert, J. Animal vectors of Arceuthobium americanum seed in Manitoba, 865

Raddi, S. see Borghetti, M., 1071

Radford, F.G. see Kalra, Y.P., 981

Radwan, M.A., Murray, M.D., and Kraft, J.M. Growth and foliar nutrient concentrations of Pacific silver fir, 1429

Real, P.L., Moore, J.A., and Newberry, J.D. Principal components analysis of tree stem profiles, 1538

Rebertus, A.J. see Veblen, T.T., 1218

Reed, D.D., and Jones, E.A. Assessing multiple estimates of plantation productivity, 948

Reich, R.M., and Arvanitis, L.G. Correcting for nonrandom

errors in a fixed-effect model, 1550

Reid, J.B. see Hallam, P.M., 480

Reid, J.B. see Hallam, P.M., 1235

Reid, M.S. see Veblen, T.T., 1218

Reiter, M.A. see Gilliam, F.S., 422

Rheney, J.W. see Oppenheimer, M.J., 329

Riitters, K.H., Stanton, B.J., and Walkup, R.H. Factor levels for density comparisons in the split-block spacing design, 1625

Roberts, B.R., and Cannon, W.N., Jr. Changes in xylem pressure potential of red spruce seedlings treated with ozone and simulated acid rain, 1200

Roberts, D.R. see Webb, D.T., 1303

Roberts, D.R. see Ellis, D.D., 1343

Roberts, L.W. see Stiff, C.M., 1330

Robinson, R.A. see McDermott, J.M., 244

Rogers, D.L., Stettler, R.F., and Heilmann, P.E. Genetic variation and productivity of *Populus trichocarpa* and its hybrids. III. Structure and pattern of variation in a 3-year field test, 372

Ross, S.D. see Bower, R.C., 1422

Rothwell, R.L. see Swanson, L.E., 1024

Roy, D.N., Konar, S.K., Banerjee, S., Charles, D.A., Thompson, D.G., and Prasad, R. Uptake and persistence of the herbicide glyphosate (Vision ®) in fruit of wild blueberry and red raspberry, 842.

Roy, G. see Gervais, C., 595

Rudolph, T.D., Kang, H.C., and Guries, R.P. Realized genetic gain for 2nd-year height in jack pine, 707

Rustad, L.E., and Cronan, C.S. Cycling of aluminum and nutrients in litterfall of a red spruce (*Picea rubens* Sarg.) stand in Maine, 18

Ryan, M.G. Sapwood volume for three subalpine conifers: predictive equations and ecological implications, 1397

Saucier, J.R. see Zahner, R., 612

Schaetzl, R.J., Johnson, D.L., Burns, S.F., and Small, T.W. Tree uprooting: review of terminology, process, and environmental implications, 1

Scherzer, A.J. see McClenahen, J.R., 880

Schiffman, P.M., and Johnson, W.C. Phytomass and detrital carbon storage during forest regrowth in the southeastern United States Piedmont, 69

Schmitz, R.F., McGregor, M.D., Amman, G.D., and Oakes, R.D. Effect of partial cutting treatments of lodgepole pine stands on the abundance and behavior of flying mountain pine beetles, 566

Schmitz, R.F. see Amman, G.D., 60

Schmitz, R.F. see Amman, G.D., 284

Schnorbus-Panozzo, E. see Weetman, G.F., 1501

Schnorbus-Panozzo, E. see Weetman, G.F., 1512

Schowalter, T.D. Canopy arthropod community structure and herbivory in old-growth and regenerating forests in western Oregon, 318

Schowalter, T.D. see Zhong, H., 943

Schreuder, H.T. see De Vries, P.G., 1067

Schuch, U.K., Duryea, M.L., and Fuchigami, L.H. Frost hardiness as acquired by Douglas-fir seedlings in three Pacific Northwest nurseries, 192

Schuch, U.K., Duryea, M.L., and Fuchigami, L.H. Dehardening and budburst of Douglas-fir seedlings raised in three Pacific Northwest nurseries, 198

Schultz, J.C. see Kleiner, K.W., 1445

Schwintzer, C.R., and Ostrofsky, A. Factors affecting germination of *Myrica gale* seeds, 1105

Seiler, J.R. see Johnsen, K.H., 1598

Semlitsch, R.D. see Weckerly, F.W., 811

Shafer, S.R., and Heagle, A.S. Growth responses of field-grown loblolly pine to chronic doses of ozone during multiple growing seasons, 821

Shiver, B.D. see Oppenheimer, M.J., 329

Shiver, B.D. see Borders, B.E., 768

Shoulders, E. see Caulfield, J.P., 743

Sidhu, S.S. see Feng, J.C., 378

Sigmon, J.T., Gilliam, F.S., and Partin, M.E. Precipitation and throughfall chemistry for a montane hardwood forest ecosystem: potential contributions from cloud water, 1240

Sigmon, J.T. see Gilliam, F.S., 422

Simon, J.-P. see Tremblay, M., 1371

Sinclair, S.A. see Cohen, D.H., 1629

Sirois, L., and Payette, S. Postfire black spruce establishment in subarctic and boreal Quebec, 1571

Small, T.W. see Schaetzl, R.J., 1

Smith, F.W. see Long, J.N., 930

Söderström, B. see Nohrstedt, H.-Ö., 323

Stanton, B.J. see Riitters, K.H., 1625

Stanturf, J.A., Stone, E.L., Jr., and McKittrick, R.C. Effects of added nitrogen on growth of hardwood trees in southern New York, 279

Stein, J. see Prévost, M., 575

Stergas, R.L., and Adams, K.B. Jack pine barrens in northeastern New York: postfire macronutrient concentrations, heat content, and understory biomass, 904

Stettler, R.F. see Rogers, D.L., 372

Stewart, W.K. see Burnes, T.A., 1642

Stiff, C.M., Wenny, D.L., Dumroese, R.K., Roberts, L.W., and Stiff, C.T. Establishment of western white pine shoots in vitro using needle fascicles, 1330

Stiff, C.T. see Stiff, C.M., 1330

Stocks, B.J. Fire behavior in mature jack pine, 783

Stocks, B.J. see Martell, D.L., 1555

Stoehr, M.U., Cai, M., and Zsuffa, L. In vitro plant regeneration via callus culture of mature Salix exigua, 1634

Stone, E.L., Jr. see Stanturf, J.A., 279

Strader, R.H., and Binkley, D. Mineralization and immobilization of soil nitrogen in two Douglas-fir stands 15 and 22 years after nitrogen fertilization, 798

Stringer, J.W., Kalisz, P.J., and Volpe, J.A. Deep tritriated water uptake and predawn xylem water potentials as indicators of vertical rooting extent in a *Quercus-Carya* forest, 627

Struve, D.K., Jett, J.B., McKeand, S.E., and Cannon, G.P. Subsoiling in a loblolly pine seed orchard: effects on seed quality, 505

Stuart, J.D., Agee, J.K., and Gara, R.I. Lodgepole pine regeneration in an old, self-perpetuating forest in south central Oregon, 1096

Sugg, D.W. see Weckerly, F.W., 811

Surles, S.E., Hamrick, J.L., and Bongarten, B.C. Allozyme variation in black locust (Robinia pseudoacacia), 471

Sutherland, J.R. see Futai, K., 1256

Swank, W.T. see Montagnini, F., 1226

Swanson, L.E., and Rothwell, R.L. Substrate freeze-thaw in a drained Alberta fen, 1024

Taylor, B.R., Parsons, W.F.J., and Parkinson, D. Decomposition of *Populus tremuloides* leaf litter accelerated by addition of *Alnus crispa* litter, 674

Taylor, S.L. see Baxter, R., 1338

Thérien, G. see Howard, A.F., 1262

Thibault, J.R. see Jobidon, R., 1291

Thier, R.W. see Amman, G.D., 60

Thompson, C. see Ying, C.C., 832

Thompson, D.C. The effect of stand structure and stand density on the leaf area – sapwood area relationship of lodgepole pine, 392

Thompson, D.G. see Roy, D.N., 842

Thompson, I.D., and Mallik, A.U. Moose browsing and allelopathic effects of Kalmia angustifolia on balsam fir regeneration in central Newfoundland, 524

Thomson, A.J., and McMinn, R.G. Height growth rates of young white spruce and lodgepole pine, 257

Thomson, A.J., and McMinn, R.G. Effects of stock type and site preparation on growth to crown closure of white spruce and lodgepole pine, 262

Thomson, T.A. Evaluating some financial uncertainties of tree improvement using the capital asset pricing model and dominance analysis, 1380

Timmer, V.R. see Munson, A.D., 162

Timmer, V.R. see Munson, A.D., 171

Tinnin, R.O. see Wanner, J.L., 736 Titus, S.J. see Johnson, A.F., 527

Toivonen, P.M.A. see Vidaver, W., 1478

Tremblay, M., and Simon, J.-P. Genetic structure of marginal populations of white spruce (*Picea glauca*) at its northern limit of distribution in Nouveau-Québec, 1371

Trevitt, A.C.F. see Beck, J.A., 791 Turnbull, C.R.A. see West, P.W., 270

Turvey, N.D. see Carlyle, J.C., 96

Valentine, D. see France, E.A., 1648 Van Cleve, K. see Dyrness, C.T., 1389

van den Driessche, R. Changes in osmotic potential of Douglasfir (Pseudotsuga menziesii) seedlings in relation to temperature and photoperiod, 413

Van Kooten, G.C., and Arthur, L.M. Assessing economic benefits of climate change on Canada's boreal forest, 463

Väre, H. The mycorrhizal condition of weakened Scots pine saplings grown on ploughed sites in northern Finland, 341

Veblen, T.T. Nothofagus regeneration in treefall gaps in northern Patogonia, 365

Veblen, T.T., Hadley, K.S., Reid, M.S., and Rebertus, A.J. Blowdown and stand development in a Colorado subalpine forest, 1218

Verbyla, D.L., and Fisher, R.F. An alternative approach to conventional soil-site regression modeling, 179

Vidaver, W., Binder, W., Brooke, R.C., Lister, G.R., and Toivonen, P.M.A. Assessment of photosynthetic activity of nursery-grown *Picea glauca* seedlings using an integrating fluorometer to monitor variable chlorophyll fluorescence, 1478

Vidaver, W.E. see Mohammed, G.H., 1322 Vimmerstedt, J.P. see McClenahen, J.R., 889

Visser, S., and Parkinson, D. Microbial respiration and biomass in soil of a lodgepole pine stand acidified with elemental sulphur,

Visser, S. see Danielson, R.M., 1412

Volpe, J.A. see Stringer, J.W., 627

von Rudloff, E., and Lapp, M.S. Some observations on the leaf oil terpene composition of mountain hemlock, *Tsuga merten*siana, 848

Waide, J.B. see Montagnini, F., 1226

Waite, C.E. see DeHayes, D.H., 1037

Walkup, R.H. see Riitters, K.H., 1625

Wang, Q. see Yang, B., 1527

Wanner, J.L., and Tinnin, R.O. Some effects of infection by Arceuthobium americanum on the population dynamics of *Pinus* contorta in Oregon, 736

Waring, R.H. see Pothier, D., 432 Waring, R.H. see Pothier, D., 1564

Wear, D.N. Structural change and factor demand in Montana's solid wood products industries, 645

Webb, D.T., Webster, F., Flinn, B.S., Roberts, D.R., and Ellis, D.D. Factors influencing the induction of embryogenic and caulogenic callus from embryos of *Picea glauca* and *P. engelmanii*, 1303

Webb, D.T. see Ellis, D.D., 1343

Webster, F. see Webb, D.T., 1303

Weckerly, F.W., Sugg, D.W., and Semlitsch, R.D. Germination success of acorns (Quercus): insect predation and tannins, 811

Weerasuriya, Y. see Norby, R.J., 889

Weetman, G.F., Fournier, R., Barker, J., Schnorbus-Panozzo, E., and Germain, A. Foliar analysis and response of fertilized chlorotic Sitka spruce plantations on salal-dominated cedar-hemlock cutovers on Vancouver Island, 1501

Weetman, G.F., Fournier, R., Barker, J., and Schnorbus-Panozzo, E. Foliar analysis and response of fertilized chlorotic western hemlock and western red cedar reproduction on salaldominated cedar-hemlock cutovers on Vancouver Island, 1512

Wenny, D.L. see Stiff, C.M., 1330

West, P.W., Beadle, C.L., and Turnbull, C.R.A. Mechanistically based, allometric models to predict tree diameter and height in even-aged monoculture of *Eucalyptus regnans* F. Muell, 270

West, R.J. Cone depredations by the red squirrel in black spruce stands in Newfoundland: implications for commercial cone collection, 1207

White, D.E., and Newton, M. Competitive interactions of whiteleaf manzanita, herbs, Douglas-fir, and ponderosa pine in southwest Oregon, 232

Whitney, R.D. Root rot damage in naturally regenerated stands of spruce and balsam fir in Ontario, 295

Wiecko, A.T. see Merkle, S.A., 285

Wiggins, K.L. see Blenis, P.V., 683

Wilkes, J. Variation in wood density of *Pinus radiata* in New South Wales, Australia, 289

Wilson, B.F. Shoot production in Douglas-fir branches, 802

Wojtkowski, P.A. see Cubbage, F.W., 759

Wooderd R.W. see Baxter, R., 1338

Woodard, P.M. see Johnson, A.F., 527

Woodcock, D.W. Climate sensitivity of wood-anatomical features in a ring-porous oak (Quercus macrocarpa), 639

Woods, J.H., and Heaman, J.C. Effect of different inbreeding levels on filled seed production in Douglas-fir, 54

Wotton, B.M. see Flannigan, M.D., 1059

Xu, L. see Kremer, R., 690

Yang, B., and Wang, Q. Distribution of the pinewood nematode in China and susceptibility of some Chinese and exotic pines to the nematode, 1527

Yaussy, D.A. Green lumber grade yields from black cherry and red maple factory grade logs sawed at band and circular mills, 1436

Yeatman, C.W. see Magnussen, S., 962

Yin, X., Perry, J.A., and Dixon, R.K. Influence of canopy removal on oak forest floor decomposition, 204

Ying, C.C., Thompson, C., and Herring, L. Geographic variation, nursery effects, and early selection in lodgepole pine, 832

Zahner, R., Saucier, J.R., and Myers, R.K. Tree-ring model interprets growth decline in natural stands of loblolly pine in the southeastern United States, 612

Zak, D.R., Host, G.E., and Pregitzer, K.S. Regional variability

in nitrogen mineralization, nitrification, and overstory biomass in northern Lower Michigan, 1521

Zarnovican, R. see Ouellet, D., 992

Zedaker, S.M. see Nicholas, N.S., 1487

Zeide, B. Accuracy of equations describing diameter growth, 1283 Zhong, H., and Schowalter, T.D. Conifer bole utilization by wood-boring beetles in western Oregon, 943

Zsuffa, L. see Stoehr, M.U., 1634